

## Maintenance Calendar for Cool-Season Turfgrass Lawns in Virginia<sup>z</sup>

*Mike Goatley, Extension specialist, Crop and Soil Environmental Sciences, Virginia Tech*

*Shawn Askew, Plant Pathology, Physiology, and Weed Science, Virginia Tech*

*Derik Cataldi and Chantel Wilson, Virginia Department of Conservation and Recreation*

Maintenance activity <sup>y</sup>	Month													
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec		
<b>Seeding<sup>x</sup></b> (Initial establishment and/or renovation)			XXXXXXXXXXXXXXXXXXXX						XXXXXXXXXXXXXXXX					
<b>N Fertilization<sup>w</sup></b>			XXXXXXXXXXXXXXXXXXXX						XXXXXXXXXXXXXXXXXXXX					xxxxx
<b>PRE herbicides<sup>v</sup></b>			XXXXXXXXXX					XXXXXXXXXX						
<b>POST herbicides<sup>u</sup></b>			XXXXXXXXXXXXXXXXXXXX						XXXXXXXXXXXXXXXXXXXX					
<b>Cultivation/dethatching</b>			XXXXXXXXXX						XXXXXXXXXXXXXXXX					

<sup>z</sup> Predominant cool-season turfgrasses for Virginia lawns are Kentucky bluegrass, tall fescue, perennial ryegrass, and fine-leaf fescues.

<sup>y</sup> Preferred timing for maintenance activity is indicated by an upper case 'X'. Second best timing indicated by lower case 'x'.

<sup>x</sup> Recommended seeding rates per 1000 sq ft are 2-3 lbs for Kentucky bluegrass; 4-6 pounds for tall fescue; 3-5 pounds for perennial ryegrass; 3-5 pounds for fine-leaf fescues. Sod is also available for most of these grasses and while these are preferred planting times, sod can be established year round as long as soils are not frozen and supplemental irrigation is available when necessary. Consult additional Virginia Cooperative Extension (VCE) publications at <http://www.pubs.ext.vt.edu/category/lawns.html> for more information.

<sup>w</sup> Applications of no more than 0.7 lb of readily available (i.e. water soluble) nitrogen /1000 sq ft per active growing month are recommend-ed during the preferred timing of fall. Use levels of 0.25-0.5 lb readily available N/1000 sq ft per growing month during secondary growing periods and never apply fertilizer to frozen soils. Consider the use of slowly available nitrogen (SAN) sources (those containing ≥ 15% water insoluble N) whenever possible (application levels of up to 0.9 lb SAN/ 1000 sq ft per active growing month) and apply other nutri-ents and/or lime based on soil test results. Note: it is recommended to test homelawn soils every 3-4 years.

<sup>v</sup> Spring preemergent (PRE) herbicide applications are primarily targeting summer annual weeds such as crabgrass, goosegrass, or foxtails. Fall applications are primarily targeting annual bluegrass and winter annual broadleaves such as henbit, deadnettle, chickweed, and gerani-um. Before applying any PRE herbicide consider possible effects it will have on seeding desirable turfgrasses in the future.

<sup>u</sup> Weeds must be actively growing to achieve control with postemergence (POST) herbicides. For cool-season weeds, treat when tempera-tures are ≥ 50° F. For warm-season weeds, temperatures ≥ 80° F are required for maximum control. Proper identification of the weed is critical in selecting appropriate control strategies. Consult VCE resources for assistance in weed identification. For chemical recommenda-tions, refer to the "Home Grounds and Animals" Pest Management Guide by VCE ([www.ext.vt.edu/pubs/pmg/](http://www.ext.vt.edu/pubs/pmg/)).

[www.ext.vt.edu](http://www.ext.vt.edu)

Produced by Communications and Marketing, College of Agriculture and Life Sciences, Virginia Tech, 2015

Virginia Cooperative Extension programs and employment are open to all, regardless of age, color, disability, gender, gender identity, gender expression, national origin, political affiliation, race, religion, sexual orientation, genetic information, veteran status, or any other basis protected by law. An equal opportunity/affirmative action employer. Issued in furtherance of Cooperative Extension work, Virginia Polytechnic Institute and State University, Virginia State University, and the U.S. Department of Agriculture cooperating. Edwin J. Jones, Director, Virginia Cooperative Extension, Virginia Tech, Blacksburg; M. Ray McKinnie, Interim Administrator, 1890 Extension Program, Virginia State University, Petersburg.

VT/0116/430-523/CSES-153NP