



Commercial Fish & Shellfish Technology Fact Sheet

Dealing with *Trichodina* and *Trichodina*-like species

Stephen A. Smith, Professor, Aquatic Medicine/Fish Health, Virginia-Maryland College of Veterinary Medicine and Michael H. Schwarz, Director, Extension Specialist, Virginia Seafood Agricultural Research and Extension Center, Virginia

What are *Trichodina* and *Trichodina*-like species?

Trichodina spp. are a group of dorsal-ventrally flattened oval ciliated protozoan parasites of marine and freshwater species of finfish. A readily distinguishable characteristic of these organisms is the presence of a prominent denticular or “tooth-like” internal cytoskeleton ring. There are three additional genera of trichodinads (*Trichodinella*, *Paratrachodina* and *Dipartiella*) which are similar in description and life cycle. While small numbers of these organisms on a fish generally do not cause much of a health problem, large numbers can cause moderate to serious pathology and ultimately, death of fish. Small fish and fry are especially susceptible, and mortality can occur quickly if undiagnosed.



Figure 1. *Trichodina* sp., note prominent “tooth-like” internal ring.

How does *Trichodina* spp. affect the fish?

Trichodina spp. cause damage by feeding on mucus and detritus covering the surface of the gills and skin of the fish causing irritation to the epithelial layer of cells. This can result in hyperplasia (proliferation) of the epithelial cells, clubbing of the gill filaments and even fusion of the gill filaments. This affects the ability of the gills to maintain optimal respiratory and excretory activities, and the ability of the skin to maintain proper homeostatic osmoregulatory properties. Massive infestations of these parasites on fish can also directly result in superficial to deep ulcerative skin lesions which then allow for secondary bacterial and fungal infections to develop at the affected site.

What is the life cycle of *Trichodina* spp.?

There are species of trichodinads that inhabit freshwater, brackish water and salt water. Some have a preference for the gill area, others for the skin, and others that occur on both the gills and



Figure 2. *Trichodina* sp. in skin scrape of fish.




Figure 3. Histopathology of *Trichodina* sp. in gill tissue of fish. Note multiple parasites in this H&E stained tissue section.

skin of fish. Trichodinads reproduce by simple binary fission under conditions that are usually optimal for the host fish. Most species are host specific and presumably spread from fish to fish by incidental contact between susceptible host fish, as well as through contact with the organism in the water column.

Why is *Trichodina* spp. a problem in aquaculture?

Trichodina spp can cause extensive fish mortality in an aquaculture system. The ability of this parasite to quickly multiply under certain environmental conditions or when the fish are stressed by other factors makes early detection of this parasite a high priority in an aquaculture facility. Once diagnosed, an appropriate treatment or management response is essential to prevent rapid loss of fish stocks.

How can *Trichodina* spp. be controlled?

There are several methods by which *Trichodina* spp. may be controlled in the aquaculture of foodfish. These include chemical treatments, freshwater baths, and flushing. UV is generally considered ineffective due to the high dosage rates required to kill the organism.

Chemical Treatment

The only FDA-approved chemical for the treatment of external parasites on foodfish is aquaculture-approved formalin. This is probably the best method to date for controlling *Trichodina* spp. infestations in an aquaculture system. A formalin bath of 170-250 ppm for 60 minutes is the FDA-approved recommendation. However, experience has shown that a single formalin bath may not completely remove all of the parasites from fish, especially marine fish, and long term or periodic treatments may be needed to keep this parasite under control. Therefore a continuous bath of 25 ppm formalin is also approved by the FDA for use on foodfish. This may affect the biofiltration of a recirculation system slightly, but usually only for a short period of time. This dosage may be repeated as necessary based upon periodic monitoring until no *Trichodina* spp. are detectable via skin scrapes. In addition, sodium chloride (salt), regarded as a Compound of Low Regulatory Priority by the FDA, may be used at 1.5-3.0 ppt to treat *Trichodina* spp. infestations on freshwater fish.

Water Bath

Another common method for controlling *Trichodina* spp. on marine finfish is to utilize periodic fresh water dips. Though stressful on fish due to increased handling and the osmotic stress, this method can be very effective in reducing the overall number of parasites on fish. This is an effective method for treating individual fish such as broodstock, but may not be a viable option in a production facility due to the logistics associated with handling and treating large numbers of fish.

Flushing

Flushing of production systems (i.e., the removal of system water prior to treatment) is another means of reducing infestation levels of *Trichodina* spp. This method may be effective by physically removing any dislodged parasites in the water column from the system.

UV Treatment

Recently, UV irradiation treatment of the water column has been examined as a potential control method for infections caused by ectoparasitic protozoans. A UV dose of $2.2 \times 10^6 \mu\text{W s/cm}^2$ of the circulating rearing water succeeded in control of *Trichodina truttae* infection in juvenile salmon (*Oncorhynchus keta*).

References

1. Gaze, W.H. and R. Wootten. 1998. Ectoparasitic species of the genus *Trichodina* (Ciliophora:Peritrichida) parasitizing British freshwater fish. *Folia Parasitologica* 45:177-190.
2. Lom, J. 1995. Protozoan and Metazoan Infections. In *Fish Diseases and Disorders*, Volume 1. P.T.K. Woo, ed. CABI Publishing, New York, NY.
3. Mizuno S., S. Urawa, M. Miyamoto, M. Hatakeyama, N. Koide and H. Ueda. 2018. Ectoparasitic protozoans *Ichthyobodo salmonis* and *Trichodina truttae* in juvenile chum salmon using ultraviolet disinfection of rearing water. *J Fish Dis.* 2018:1–12.
4. Smith, S.A. and E.J. Noga. 1992. General Parasitology. In *Fish Medicine*. M.K. Stoskopf, ed. W.B. Saunders Co. Philadelphia, PA.
5. Wellborn, T.L. 1967. *Trichodina* (Ciliata: Urceolariidae) of Freshwater Fishes of the Southeastern United States. *J. Protozool.* 14:399-412.

This fact sheet was published by:
Virginia Tech, the Virginia Sea Grant College Program, and Virginia Cooperative Extension

Photos: Stephen Smith, DVM, PhD

For further information contact:

Stephen A. Smith

Virginia-Maryland College of Veterinary Medicine

stsmith7@vt.edu

This publication is the result of research supported by the NOAA Office of Sea Grant, the U.S. Department of Commerce, and the Virginia Sea Grant College Program.



VSG-18-5

