



# ***Trueperella pyogenes*: A Practical Summary for Controlling Mastitis**

*Authored by Christina S. Petersson-Wolfe, Associate Professor & Extension Dairy Scientist, School of Animal Science, Virginia Tech; Turner Swartz, former Ph.D. student, Dairy Science, Virginia Tech; and Sebastián Umaña Sedó, Assistant Professor, Virginia-Maryland College of Veterinary Medicine*

*Trueperella pyogenes* are environmental pathogens that typically cause an acute, purulent form of mastitis that is sometimes referred to as “summer mastitis.” *Trueperella pyogenes* are Gram-positive bacteria that can appear like coccobacilli — an intermediate between cocci and bacilli. When plated on blood agar, scant to no growth is visible at 24 hours. However, at 48 hours, tiny, pinpoint, smooth, whitish colonies develop and are surrounded by a narrow zone of hemolysis.

Information in this publication was summarized from the National Mastitis Council’s Laboratory Handbook on Bovine Mastitis (Hogan et al. 1999).

## **Where are these organisms found?**

Reservoirs of *Trueperella pyogenes* include teat injuries, wound infections, udder infections, abscesses, and genital tracts.

## **How do *Trueperella pyogenes* spread to the mammary gland?**

*Trueperella pyogenes* is spread by flies, though recent research shows its transmission cannot be fully attributed to flies alone (Wente et. al., 2024).

## **How can you prevent and control mastitis caused by *Trueperella pyogenes*?**

Prevention must include an effective fly control program as well as maintaining clean, dry environments, particularly for calving areas. Fly control programs

should include routine manure removal; maintaining clean, dry pens; routine cleaning of waterers and feed bunk areas; and the use of insecticides and/or fly traps.

Other control strategies include “killing” or drying off the infected quarter or removing the cow from the herd.

## **When are *Trueperella pyogenes* mastitis infections most likely to occur?**

*Trueperella pyogenes* typically cause mastitis in dry cows or heifers prior to calving, particularly during the summer or humid weather. It is for this reason it is sometimes referred to as “summer mastitis.” Infection can occur in lactating animals with teat injuries or due to the use of cannulas or dilators.

## **How likely is *Trueperella pyogenes* mastitis to cure?**

For new infections detected early and confirmed to be caused by *Trueperella pyogenes*, treatment may be considered using labeled intramammary (IMM) therapy products or a 4–5 day course of an injectable broad-spectrum antimicrobial. Producers should always consult with their herd veterinarian before initiating any treatment protocol.

Once the infection becomes established, the prognosis is generally poor, and affected cows will often lose function of the infected quarter. To help prevent the spread of *Trueperella pyogenes* within the herd, it is typically recommended that the affected quarter be dried off or that the cow be removed from the herd.

## Quick Notes

- *Trueperella pyogenes* are environmental pathogens associated with teat injuries.
- *Trueperella pyogenes* mastitis typically occurs prior to calving, during the dry period.
- *Trueperella pyogenes* are spread by flies; thus, an effective fly control program is very important.
- Prognosis after infection is established is poor; therefore, “killing” or drying of the quarter or removing the cow from the herd is necessary to prevent the spread of this pathogen.

## References

- Hogan, J. S., R. N. Gonzalez, R. J. Harmon, S. C. Nickerson, S. P. Oliver, J. W. Pankey, and K. L. Smith. (1999). *Laboratory Handbook on Bovine Mastitis*. Madison, WI: National Mastitis Council.
- Wente, N., Leimbach, S., Woudstra, S., & Krömker, V. *Trueperella Pyogenes—Strain Diversity and Occurrence in Dairy Herds*. *Pathogens*, 13(7), 534. 2024.

---

Visit our website: [www.ext.vt.edu](http://www.ext.vt.edu)

Produced by Virginia Cooperative Extension, Virginia Tech, 2026

Virginia Cooperative Extension is a partnership of Virginia Tech, Virginia State University, the U.S. Department of Agriculture (USDA), and local governments, and is an equal opportunity employer. For the full non-discrimination statement, please visit [ext.vt.edu/accessibility](http://ext.vt.edu/accessibility).

VT/0626/DASC-71P (DASC-208P)